

M.Sc. PHYSICS
FIRST SEMESTER
STATISTICAL PHYSICS
MSP – 102 [REPEAT]
[USE OMR FOR OBJECTIVE PART]

**SET
A**

Duration: 3 hrs.

Full Marks: 70

Time: 30 min.

(Objective)

Marks: 20

1 × 20 = 20

Choose the correct answer from the following:

- In canonical ensemble, the r.m.s fluctuation in energy is
 - $\frac{\sqrt{kT^2}}{U}$
 - $\sqrt{kTC_v}$
 - $\sqrt{\frac{kT^2 C_v}{U}}$
 - $\frac{U}{\sqrt{kT^2 C_v}}$
- Which of the following statement is false?
 - In classical statistics, the particles have a certain degree of togetherness as well as separateness.
 - Maxwell-Boltzmann statistics describes the distribution of gas molecules.
 - Liouville's theorem gives the principle of conservation of energies of particles.
 - In Grand canonical ensemble, the system is separated by rigid, permeable and conducting walls.
- In case of Maxwell-Boltzmann velocity distribution curve, which one of the following is correct?
 - As T increases, the distribution becomes narrow.
 - As T increases, the distribution spreads out.
 - As T increases, the distribution gets sharper.
 - As T decreases, the distribution spreads out.
- In case of Maxwell-Boltzmann statistics, the molecular size is
 - Negligible
 - Equal to the intermolecular distance
 - Less than the intermolecular distance
 - More than the intermolecular distance
- The thermodynamic probability of Maxwell-Boltzmann distribution is
 - $N! \frac{g_i^{n_i}}{n_i!}$
 - $N! \frac{g_i^{n_i}}{n_i}$
 - $N \frac{g_i^{n_i}}{n_i!}$
 - $N! \frac{g_i^{n_i}}{n_i!}$
- The partition function for a 3 dimensional monoatomic gas is given by?
 - $\frac{1}{h^3} \iiint e^{-E/kT} d^3q d^3p$
 - $\frac{1}{h} \iint e^{-\beta E} d^3q d^3p$
 - $\frac{1}{h^2} \iint e^{-E/kT} dq dp$
 - $\frac{1}{h^2} \iint e^{-E/kT} d^3q d^3p$

7. In case of a classical ideal gas, which of the following option for N particle system is correct?
- a. $Z_N = \left(\frac{V(\sqrt{2\pi mkT})^3}{h^3} \right)^N$
- b. $Z_N = \left(\frac{V2\pi mkT}{h^3} \right)^N$
- c. $Z_N = \left(\frac{V(2\pi mkT)^3}{h^3} \right)^N$
- d. $Z_N = \left(\frac{V\sqrt{2\pi mkT}}{h^3} \right)^N$
8. Maxwell-Boltzmann distribution function is given by $n_i =$
- a. $\frac{g_i}{e^{\alpha+\beta E_i}-1}$
- b. $\frac{g_i}{e^{\alpha+\beta E_i+1}}$
- c. $\frac{g_i}{e^{\alpha-\beta E_i}}$
- d. $\frac{g_i}{e^{\alpha+\beta E_i}}$
9. In canonical ensemble, which of the following is true?
- a. Energy can vary from 0 to infinity
- b. Energy does not vary at all
- c. Energy can vary from 0 to 1
- d. Energy is restricted
10. Partition function of an equilibrium system is given by
- a. $\sum_i g_i e^{-\beta E_i}$
- b. $\sum_i g_i e^{-\beta E_i}$
- c. $\sum_i e^{-\beta E_i}$
- d. $\sum_i g_i e^{\alpha+\beta E_i}$
11. What is the partition function in quantum statistical mechanics?
- a. It represents the total number of accessible microstates for a given macrostate
- b. It represents the probability of a system being in a particular energy state
- c. It represents the total energy of a system
- d. It represents the average value of an observable in a given quantum state
12. Which of the following is a fundamental postulate of quantum statistical mechanics?
- a. The wave function collapses to an eigenstate upon measurement
- b. The total energy of a system is conserved
- c. The probability of an event is given by the square of the absolute value of the wave function
- d. The average value of an observable is given by the expected value of the corresponding operator
13. At what temperature does Bose-Einstein condensation occur?
- a. Room temperature
- b. Absolute zero
- c. 1000 degrees Celsius
- d. It can occur at any temperature
14. What are some real-world applications of Bose-Einstein statistics?
- a. Superconductivity and superfluidity
- b. Solar energy capture
- c. Nuclear fission
- d. Quantum computing
15. In quantum statistical mechanics, the average energy of a system is given by
- a. The expectation value of the Hamiltonian operator.
- b. The product of the partition function and the temperature
- c. The sum of the energies of all possible states divided by the total number of states
- d. The Gibbs free energy

16. The grand canonical density matrix can be written as (symbols have usual meaning)
- | | |
|-------------------------------------|--|
| a. $\rho_{nn} = e^{+\beta E_n} / Z$ | b. $\rho_{nn} = e^{-\beta(E_i - \mu N_i)} / Z$ |
| c. $\rho_{nn} = e^{E_n} / Z$ | d. $\rho_{nn} = e^{\beta(E_n - n_i)} / Z$ |
17. In Fermi-Dirac distribution function, the chemical potential represents
- | | |
|--|--|
| a. The rate of change of particle number with respect to energy. | b. The energy required to add one more particle to the system. |
| c. The temperature of the system. | d. The probability of finding a particle in a specific energy state. |
18. In a quantum mechanical ensemble, which of the following best describes the behavior of individual particles?
- | | |
|--|--|
| a. The behavior of individual particles can be predicted with certainty. | b. The behavior of individual particles follows classical laws of physics. |
| c. The behavior of individual particles is probabilistic. | d. The behavior of individual particles is completely random. |
19. What is the significance of Pauli exclusion principle in quantum statistical mechanics?
- | | |
|--|--|
| a. Determines the probability distribution of particles in different energy states | b. Ensures the stability of electrons in atoms and molecules |
| c. Describes the behavior of identical particles in quantum systems | d. Determines the average energy of a system |
20. Number of microstates in a macrostate may be
- | | |
|-----------|------------|
| a. Equal | b. \geq |
| c. \leq | d. greater |

--- --

(Descriptive)

Time : 2 hrs. 30 min.

Marks : 50

[Answer question no.1 & any four (4) from the rest]

- | | | | |
|----|----|--|----------|
| 1. | a. | Discuss briefly the phase space of a classical system and derive the partition function of classical ideal gas for N particle system. | 5+5=10 |
| | b. | Explain Bose-Einstein condensation and derive the criteria for Bose Einstein condensation to occur. | |
| 2. | | Find out the energy and density fluctuation in grand canonical ensemble. | 4+6=10 |
| 3. | | Discuss the Simple Harmonic Oscillator in classical statistics along with three thermodynamic properties. | 8+2=10 |
| 4. | | Derive Maxwell-Boltzmann velocity distribution function and draw the graph showing the dependence on temperature. | 6+4=10 |
| 5. | a. | Explain the statistical weight of BE statistics. | 2+8=10 |
| | b. | Derive the equation of state for ideal Bose gas and discuss classical limit for Bosons. | |
| 6. | a. | Find the expression for average occupancy of single particle energy state in the case of Bose-Einstein and Fermi-Dirac distribution assuming ideal gas conditions. | 6+2+2=10 |
| | b. | Derive Liouville's equation (time evolution of density matrix) in context of quantum statistical mechanic | |
| | c. | Write a brief note on quantum mechanical microcanonical ensemble. | |
| 7. | a. | What do you mean by ideal Fermi gas? Derive the expression for Fermi energy. | 1+4=5 |
| | b. | Show graphically how the distribution functions (Maxwell-Boltzmann, Bose-Einstein and Fermi Dirac) varies with temperature $((\epsilon - \mu)/KT)$. | 1 |
| | c. | What is Ising model? Provide a concise explanation of how the Ising model is used to simulate the physics of a ferromagnetic substance. | 3 |
| 8. | a. | What is density matrix? What is the role of density matrices in quantum statistical mechanics? State the properties of density matrix. | 1+1+2=4 |
| | b. | What is phase transition? Draw the phase diagram | 2+1=3 |
| | c. | State the properties of liquid Helium II. | 3 |

== *** ==