B.Sc. PHYSICS FIRST SEMESTER PHYSICS-I

BSP-711 [REPEAT] IUSE OMR FOR OBJECTIVE PARTI

Duration: 3 hrs.

Full Marks: 70

Time: 30 min.

c. Viscosity of liquid

a. $M^{-1}L^2T^{-2}$

c. M112T-2

8. The dimension of gravitational constant G is

Objective)

Marks: 20

1X20=20 Choose the correct answer from the following: Curl grad F = ? a. 0 b. 1 If a vector field $\vec{F} = x\hat{i} + 2y\hat{j} + 3z\hat{k}$, then $\nabla \times \nabla \times \vec{F}$ is a. 0 d. $3\hat{k}$ c. 2i 3. The work done of vectors force F and distance d, separated by angle θ can be calculated using a. Cross product b. Dot product c. Vector addition d. Vector subtraction 4. Pick the wrong answer a. The dot product of two vectors is a b. The cross product of two vectors is a scalar d. Curl of gradient of a vector is infinity c. Addition of vector is commutative. 5. The divergence of position vector is b. 1 a. 0 c. 2 d. 3 6. If the surface of a liquid is plane, then the angle of contact of the liquid with the walls of container is a. Acute angle b. Obtuse angle c. 900 d. 00 7. When there are no external forces, the shape of a liquid drop is determined by a. Surface tension of the liquid b. Density of liquid

d. Temperature of air only

b. $M^{-1}L^3T^{-2}$

d. M-1L2T-1

| | Artinertial frame of reference is that frame a. in which the Newton's 1st law is valid c. which is not moving with uniform velocity | b. in which law of inertia is valid d. in which both 1st law of motion and law of inertia is valid |
|----------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------|
| | 10. Earth isa. A non-inertial framec. Both of a. and b. | b. An inertial frame d. Nonc of a. and b. |
| | 11. Moment of inertia of a solid sphere is a. $\frac{2}{3}M^2r^2$ c. $\frac{2}{5}Mr^2$ | b. $\frac{3}{2}Mr^2$ d. $\frac{5}{2}Mr^2$ |
| | 12. The surface of water in contact with glass wa a. Plane c. Convex | all is b. Concave d. Both (a) and (b) |
| | 13. The calculation of the moment of the body d a. Moment c. Moment of Inertia | ue to the loadings involve a quantity called b. Inertia d. Rotation |
| | 14. The axis about which moment of area is takea. Axis of areac. Axis of reference | n is known as b. Axis of moment d. Axis of rotation |
| | 15. The condition for conservative force is a. $F = \nabla . U$ c. $F = -\nabla U$ | b. $F = -\nabla . U$ d. $F = -\nabla^2 . U$ |
| | 16. According to Kepler, the line joining a plane equal interval of time. (Fill the blank) a. Equal area c. One third of area | b. Equal length d. None of these |
| | 17. Example of an simple harmonic motion (SHM) isa. Vibration of a springc. None of a. and b. | b. Motion of a pendulum d. Both of a. and b |
| 18. The maximum displacement of a vibrating particle is called its a. Velocity b. Amplitude c. Acceleration d. Phase | | b. Amplitude |
| | | |

19. The time period of a vibrating particle is a.
$$T = 2\pi \sqrt{\frac{Displacement}{Acceleration}}$$
 c.
$$T = \sqrt{\frac{Displacement}{Acceleration}}$$

$$T = \sqrt{\frac{Displacement}{Acceleration}}$$

b.
$$T = 2\pi \sqrt{\frac{Acceleration}{Displacement}}$$

d.
$$T = 2\pi \sqrt{\frac{Displacement^2}{Acceleration}}$$

20. A vector point A vertically upward and point B towards north. The vector product $A \times B$

a. along west c. zero

b. along east

d. vertically downward

3

(<u>Descriptive</u>)

Time: 2 hrs. 30 min.

Marks: 50

[Answer question no.1 & any four (4) from the rest]

- 1. Calculate the minimum velocity and period of revolution of an artificial satellite at a height *h* from the surface of the earth. A small artificial satellite is revolving round the earth very close to it. If the radius of the earth is about 6400km and acceleration due to gravity 9.8 m/sec², find the period of revolution of the satellite.
- 2. (i) Define escape velocity.

1+3+2+4

=10

- (ii) Derive an expression for escape velocity in terms of acceleration due to gravity and radius of the earth.
- (iii) Calculate the value of escape velocity if the radius of earth is 6.4×10^6 m and g = 9.8 ms⁻².
- (iv) Moon has no atmosphere. Why?
- 3. (i) Define moment of inertia.

1+3+3+3

- (ii) Calculate the moment of inertia of a rectangular lamina (a) when the axis passing through the centre and perpendicular to its length and (b) when axis passing through the centre and perpendicular to the plane of lamina.
- (iii) Assuming earth to be a sphere of uniform density 5520 kg.m⁻³ and radius 6400 km, calculate the moment of inertia about its axis.
- 4. (i) Define angle of contact.

5.

2+8=10

- (ii) Show that the height of liquid risen in a capillary tube varies inversely as the radius of the tube.
- (i) Write a short note on (a) Surface tension, (b) surface energy and capillarity.

(ii) A soap film is formed on a rectangular frame 7cm dipping into the soap solution. The frame hangs from the arms of a balance. An extra weight of 0.38g is to be placed in the opposite pan to balance the pull on the frame. Calculate the surface tension of the soap solution.

4

USTM/COE/R-01

6. (i) State Gauss's divergence theorem.

- 2+4+4
- (ii) If $\vec{A} = (3x^2 + 6y)\hat{i} 14yz\hat{j} + 20xz^2\hat{k}$, evaluate the line integral $\oint \vec{A} \cdot \vec{dr}$ from (0,0,0) to (1,1,1) along the curve C. $x = t, y = t^2, z = t^3$.
- (iii) If $\vec{F} = 2z\hat{i} x\hat{j} + y\hat{k}$, evaluate $\iiint_{\vec{F}} \vec{F} dv$ where, v is the region bounded by x=0, x=2, y=0, y=4, z=x², z=2.
- 7. Define divergence of a vector function. When a vector function is said to be solenoidal. If **a** is a constant vector then prove that $\nabla \cdot (\mathbf{a} \times \mathbf{r}) = 0$. Also find the value of $\nabla \cdot (\frac{a \times r}{|r|^n})$, where $\mathbf{r} = x\hat{i} + y\hat{j} + z\hat{k}$
- 8. Explain simple harmonic motion. Show that for a body vibrating simple harmonically the time period is given by $t=2\pi\sqrt{\frac{displacement}{acceleration}}$. Represent Simple harmonic motion graphically.
